



ASPIRE
GALDERMA REWARDS



Get Rewarded for
Looking Your Best*

[ASPIREwards.com](https://www.ASPIREwards.com)

*Terms and conditions apply. See details at www.ASPIREwards.com.



Discover the difference with ASPIRE Galderma Rewards

It's the rewards program you've been waiting for. You'll get savings on all the products you love as well as gift cards, treatment certificates, and many more opportunities to earn valuable rewards. To make you look and feel your best, we connect you with aesthetic specialists and treatment details that will guide you through your journey every step of the way.

Start your aesthetic journey right now and receive a **\$20 treatment certificate** just for signing up!

Restylane

Sculptra
aesthetic

Dysport
abobotulinumtoxinA[‡]
300 Units for Injection

[‡]Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 10-13.

Join now by going online at
[ASPIRErewards.com](https://www.aspiregalderma.com)



Earn valuable rewards for every treatment you receive

For every Galderma aesthetic treatment you receive while in the program, you'll earn treatment points that can be redeemed for valuable rewards.

Treatment points†

Restylane® Family of Products = **up to 400 points**/visit, depending on treatment amount

Sculptra® Aesthetic = **250 points**/vial up to 500 points/visit

Dysport®† (abobotulinumtoxinA) = **200 points**/visit
300 Units for Injection

†Limit of points to be earned per day: 1,000 points. Point expiration: 12 months from date earned (6 months from date issued for treatment certificates)

‡Please see full Important Safety Information, including Distant Spread of Toxin Effect Boxed Warning, on pages 10-13.

There are many ways to earn bonus points

ASPIRE Galderma Rewards members can earn by:



taking our monthly quizzes



participating in surveys



referring friends

There's no paperwork required to enroll and you can manage your rewards right on your smartphone, anytime!

**Sign up now at
ASPIRErewards.com!**



A rewarding experience at every level

ASPIRE Galderma Rewards has 5 levels of loyalty.

“My positive experiences with ASPIRE Galderma Rewards began at the initial registration and have remained not only positive, but outstanding! I am grateful for the very generous discounts, treatment certificates, and fun perks. The ASPIRE Rewards program is a true gift.”

—Stephanie
Danville, CA

Your ASPIRE Galderma Rewards membership and privileges continue to improve along with your status. Points are earned based on the calendar year. The number of points you earn determines your program level. You will retain your program level while you are a member, as long as you purchase at least 1 Galderma aesthetic treatment each year. If you do not purchase at least 1 Galderma aesthetic treatment during a year, your membership will reset to Base level on January 1 of the following year and you will keep your points, unless they expire or are forfeited per the program terms and conditions.

Program level	Treatment points required	What's in it for you?
VIP	1,500+	Enjoy elite status with the greatest access to rewards, including an additional \$100 treatment certificate, and the ability to redeem exclusive VIP experiences
Royalty	1,000-1,499	Unlock even more rewarding experiences like instant \$100 treatment certificates and retailer rewards. Refer your friends to ASPIRE and you all receive valuable rewards
Insider	500-999	Start turning your points into cash via check cards that you can use however you want, whenever you want
Connector	250-499	Get even better rewards including aesthetic treatment certificates and reward cards from partner retailers like Nordstrom and Sephora
Base	0-249	Sign up for your free membership to earn points right away and gain instant access to insider info and rewards

Important Safety Information for *Dysport*[®]

Dysport is a prescription injection for temporary improvement in the look of moderate to severe frown lines between the eyebrows (glabellar lines) in adults less than 65 years of age.

What is the most important information you should know about *Dysport*?

Spread of Toxin Effects: In some cases, the effects of *Dysport* and all botulinum toxin products may affect areas of the body away from the injection site. These effects can cause symptoms of a serious condition called botulism. Symptoms of botulism can happen hours to weeks after injection and may include swallowing and breathing problems, loss of strength and muscle weakness all over the body, double vision, blurred vision and drooping eyelids, hoarseness or change or loss of voice, trouble saying words clearly, or loss of bladder control. Swallowing and breathing problems can be life threatening and there have been reports of death.

The risk of symptoms is probably greatest in children treated for muscle spasms but symptoms can also occur in adults treated for muscle spasms and other conditions, particularly in those patients who have underlying conditions that would predispose them to these symptoms.

The toxic effects have been reported at doses similar to those used to treat muscle spasms in the neck. Lower doses, in both approved and unapproved uses, have also caused toxic effects. This includes treatment of children and adults for muscle spasms.

These effects could make it unsafe for you to drive a car, operate machinery, or do other dangerous activities.

Do not have *Dysport* treatment if you: are allergic to *Dysport* or any of its ingredients (see the end of the Medication Guide for a list of ingredients), are allergic to cow's milk protein, had an allergic reaction to any other botulinum toxin product, such as Myobloc[®] or Botox[®], or have a skin infection at the planned injection site.

The dose of *Dysport* is not the same as the dose of any other botulinum toxin product. The dose of *Dysport* cannot be compared to the dose of any other botulinum toxin product you may have used.

***Dysport* may not be right for you if:** you have surgical changes to your face, very weak muscles in the treatment area, your face looks very different from side to side, the injection site is inflamed, you have droopy eyelids or sagging eyelid folds, deep facial scars, thick oily skin, or if your wrinkles can't be smoothed by spreading them apart.

Tell your doctor about all your medical conditions, including if you have: a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis, or Lambert-Eaton syndrome), allergies to any botulinum toxin product or had any side effect from any botulinum toxin product in the past, a breathing problem (such as asthma or emphysema), swallowing problems, bleeding problems, diabetes, or a slow heart beat or other problem with your heart rate or rhythm, plans to have surgery, had surgery on your face, weakness of your forehead muscles (such as trouble raising your eyebrows), drooping eyelids, or any other change in the way your face normally looks. Patients with a disease that affects muscles and nerves who are treated with typical doses of *Dysport* may have a higher risk of serious side effects, including severe swallowing and breathing problems.

Human Albumin

This product contains albumin taken from human plasma. Steps taken during donor screening and product manufacturing processes make the risk of spreading viral diseases extremely rare. In theory, there is also an extremely rare risk of contracting Creutzfeldt-Jakob disease (CJD). No cases of spread of viral diseases or CJD have ever been reported for albumin.

Allergic Reaction to Injecting in the Skin

It is not known if an allergic reaction can be caused by injecting *Dysport* into the skin. The safety of treating excessive sweating with *Dysport* is not known.

Common Side Effects

The most common side effects are nose and throat irritation, headache, injection site pain, injection site skin reaction, upper respiratory tract infection, eyelid swelling, eyelid drooping, sinus inflammation, and nausea.

Tell your doctor about all the medicines you take, including prescription and nonprescription medicines, vitamins and herbal and other natural products. Using *Dysport* with certain other medicines may cause serious side effects. **Do not start any new medicines while taking *Dysport* without talking to your doctor first.**

Especially tell your doctor if you: have received any other botulinum toxin product in the last four months, have received injections of botulinum toxin, such as Myobloc® (rimabotulinumtoxinB) or Botox® (onabotulinumtoxinA) in the past (be sure your doctor knows exactly which product you received), have recently received an antibiotic by injection, take muscle relaxants, take an allergy or cold medicine, or take a sleep medicine.

Use in Specific Populations

Dysport should not be used in children or in women who are pregnant or breastfeeding.

Ask your doctor if *Dysport* is right for you.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see *Dysport* Full Prescribing Information including Medication Guide.

Important Safety Information for the *Restylane*® Family Of Products

Indications: The *Restylane* family of products includes *Restylane*®, *Restylane-L*®, *Restylane*® *Lyft with Lidocaine* and *Restylane*® *Silk*. *Restylane*, *Restylane-L*, and *Restylane Lyft with Lidocaine* are indicated for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds.

Restylane and *Restylane-L* are indicated for mid-to-deep dermal implantation. *Restylane* and *Restylane-L* are also indicated for submucosal implantation for lip augmentation in patients over the age of 21.

Restylane Lyft with Lidocaine is indicated for implantation into the deep dermis to superficial subcutis. *Restylane Lyft with Lidocaine* is also indicated for cheek augmentation and for the correction of age-related midface contour deficiencies in patients over the age of 21.

Restylane Silk is indicated for submucosal implantation for lip augmentation and dermal implantation for correction of perioral rhytids in patients over the age of 21.

Products in the *Restylane* family contain traces of gram-positive bacterial protein and are contraindicated for patients with allergies to such material or in patients with severe allergies that have required in-hospital treatment. These products should not be used by patients with bleeding disorders or by pregnant or breastfeeding women. *Restylane-L*, *Restylane Silk* and *Restylane Lyft with Lidocaine* should not be used by anyone with a known allergy to lidocaine. Products should not be injected anywhere except areas indicated in their Instructions for Use.

Use of products in the *Restylane* family at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. The most commonly observed side effects are swelling, redness, pain, bruising, headache, tenderness, and itching at the injection site. These are typically mild in severity and typically resolve in less than 7 days in nasolabial folds and less than 14 days in lips. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis at the injection site. Do not implant into blood vessels. Use with caution in patients recently treated with anticoagulant or platelet inhibitors to avoid bleeding and bruising.

The *Restylane* family of products is available only through a licensed practitioner. Complete Instructions for Use are available at www.RestylaneUSA.com.

Important Safety Information for *Sculptra*® Aesthetic

Indications: *Sculptra Aesthetic (injectable poly-L-lactic acid)* is indicated for use in people with healthy immune systems as a single regimen for the correction of shallow to deep nasolabial fold contour deficiencies and other facial wrinkles in which deep dermal grid pattern (cross-hatch) injection technique is appropriate.

Sculptra Aesthetic should not be used by people that are allergic to any ingredient of the product or have a history of keloid formation or hypertrophic scarring. Safety has not been established in patients who are pregnant, lactating, breastfeeding, or under 18 years of age.

Sculptra Aesthetic has unique injection requirements and should only be used by a trained physician. Contour deficiencies should not be overcorrected because they are expected to gradually improve after treatment.

Sculptra Aesthetic should not be injected into the blood vessels as it may cause vascular occlusion, infarction or embolic phenomena. Use at the site of skin sores, cysts, pimples, rashes, hives or infection should be postponed until healing is complete. *Sculptra Aesthetic* should not be injected into the red area (vermillion) of the lip or in the peri-orbital area.

The most common side effects after initial treatment include injection site swelling, tenderness, redness, pain, bruising, bleeding, itching and lumps. Other side effects may include small lumps under the skin that are sometimes noticeable when pressing on the treated area. Larger lumps, some with delayed onset with or without inflammation or skin discoloration, have also been reported.

Sculptra Aesthetic is available only through a licensed practitioner. Complete Instructions for Use are available at www.SculptraAesthetic.com.



Start your
aesthetic journey
today!

Join now at ASPIRErewards.com

Please call 1-844-5ASPIRE if you have
any questions.



©2016 Galderma Laboratories, L.P.
The *Dysport* trademark is used under license.
All trademarks are the property of their respective owners.
MP-ASP-00112 03/16